## 4.6. Prevalence of Selected Chronic Conditions By Age and Gender - Per 1,000 Persons<sup>1,2</sup> in Hawai`i, Hawai`i Health Survey (HHS) 2000

Demographic Characteristic	Age in Years									Total <sup>3</sup>	
	0-14	15-17	18-24	25-34	35-44	45-54	55-64	65-74	75+	N	Prevalence/ per 1,000 Persons
Condition Type					Female						
Arthritis	0.5*	1.4*	10.5	15.4	32.9	100.8	187.5	290.4	374.2	53,115	90.5
Asthma	98.9	156.5	77.9	83.8	77.2	86.1	81.2	71.4	66.6	50,422	85.9
Diabetes	0.4*		3.7	20.4	39.2	54.2	114.2	161.8	139.8	29,798	50.8
High Blood Cholesterol	4.4	2.8*	12.9	40.2	88.0	158.5	323.0	394.5	362.1	77,614	132.2
Hypertension	1.9*	7.5*	20.7	31.8	86.4	169.5	308.7	456.8	468.6	84,569	144.1
Total	120,494	21,932	53,038	67,666	93,476	85,116	56,617	47,559	41,017	586,915	
Condition Type					Male						
Arthritis	0.3*	2.8*	4.6*	16.4	22.8	69.3	109.8	169.1	267.0	29,763	52.3
Asthma	148.2	184.1	96.3	61.2	58.4	73.6	36.0	48.8	59.5	49,569	87.1
Diabetes			3.3	9.5	25.1	52.2	124.5	153.4	115.9	23,313	41.0
High Blood Cholesterol	4.3	5.3*	14.6	53.5	133.5	235.6	333.1	333.8	294.1	76,277	134.0
Hypertension	3.6	9.8*	20.0	70.6	115.7	225.9	338.3	419.3	400.2	82,076	144.2
Total	127,337	23,339	50,618	68,339	95,509	84,941	49,273	38,657	31,086	569,099	1,156,014

<sup>&</sup>lt;sup>1</sup> All household members

Sample size 17,183 people

<sup>&</sup>lt;sup>2</sup> Sample numbers provisionally weighted and adjusted\* for total population of Hawai`i

<sup>\*(</sup>total numbers are adjusted as Ni`ihau, Kalawao, group quarters, and homeless are not represented, see HHS Procedure Manual 2000)

<sup>&</sup>lt;sup>3</sup> Row and/or column totals may not sum to totals listed due to rounding

Number of cases in sample numerator or denominator too small for reliability - shown for consistency only

<sup>--</sup> No reported cases